

# *Angelica pyrenaea*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **4**

Moisture indicator value: **8**

Reaction indicator value: **2.5**

Nutrient indicator value: **1.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.51**

Disturbance frequency (herb layer): **1.64**

Disturbance severity: **0.33**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.23**

Grazing pressure: **0.23**

Soil disturbance: **0.16**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CH \(ULI\) \*Calluno-Ulicetea\*](#), [CI \(NAR\) \*Nardetea strictae\*](#), [PA \(SCH\) \*Scheuchzerio palustris-Caricetea fuscae\*](#)

Broad habitat

Occurrence in broad habitats: **Mire, Grassland (non-alpine, non-saline), Heathland**