Anisantha diandra

Habitus and growth type

Plant height [m]: **0.43**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **16.75**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **8.44**
Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**
Moisture indicator value: **3.2**
Reaction indicator value: **6.8**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.05**
Disturbance frequency (herb layer): **1.76**
Disturbance severity: **0.77**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.71**
Mowing frequency: **0.55**
Grazing pressure: **0.2**
Soil disturbance: **0.44**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FJ (TRA) *Stipo-Trachynietea distachyae*, QC (CHE) *Chenopodietea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **V32 Mediterranean subnitrophilous annual grassland**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz