

Anisantha fasciculata

Habitus and growth type

Plant height [m]: **0.15**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **18.63**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **11**

Moisture indicator value: **2**

Reaction indicator value: **7**

Nutrient indicator value: **3**

Salinity indicator value: **1**

Disturbance indicator values

Disturbance frequency: **0.85**

Disturbance frequency (herb layer): **1.72**

Disturbance severity: **0.76**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **0.13**

Grazing pressure: **0.35**

Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FJ \(TRA\) *Stipo-Trachynietea distachyae*](#), [QC \(CHE\) *Chenopodietea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S63 Eastern garrigue](#), [S72 Eastern Mediterranean spiny heath \(phrygana\)](#)

Constant species of EUNIS habitats: [S63 Eastern garrigue](#), [S72 Eastern Mediterranean spiny heath \(phrygana\)](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**