Anisantha rubens

Habitus and growth type

Plant height [m]: **0.26**
Life span: **Annual**
Life form: **Therophyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **15.66**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.67**
Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **9.2**
Moisture indicator value: **1.8**
Reaction indicator value: **7.3**
Salinity indicator value: **1**

Disturbance indicator values

Disturbance frequency: **1.24**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.61**
Disturbance severity: **0.74**
Disturbance severity (herb layer): **0.7**
Mowing frequency: **0.37**
Grazing pressure: **0.27**
Soil disturbance: **0.38**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S66 Mediterranean halo-nitrophilous scrub**
Constant species of EUNIS habitats: **S66 Mediterranean halo-nitrophilous scrub**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz