Anthericum liliago

Habitus and growth type

Plant height [m]: **0.43**
Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **18.05**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **6.44**
Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.2**
Temperature indicator value: **6.1**
Moisture indicator value: **3.1**
Reaction indicator value: **6.6**
Nutrient indicator value: **2.8**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.5**
Disturbance severity: **0.55**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.19**
Grazing pressure: **0.28**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***, **FP (ONO) *Festuco hystricis-Ononidetea striatae***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Heathland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **4**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz