

Anthriscus cerefolium

Habitus and growth type

Plant height [m]: **0.44**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **52.68**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **2.27**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.2**

Temperature indicator value: **6.7**

Moisture indicator value: **5.1**

Reaction indicator value: **6.7**

Nutrient indicator value: **8.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.2**

Disturbance frequency (herb layer): **1.37**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.12**
Grazing pressure: **0.21**
Soil disturbance: **0.18**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QG \(EPI\) *Epilobietea angustifolii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [N1E Black Sea broad-leaved coastal dune forest](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: 7
Continentality amplitude: 5