

Anthyllis cytisoides

Habitus and growth type

Plant height [m]: **0.8**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **8.31**

Fruit, seed and dispersal

Seed mass [mg]: **2.8**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **8**

Moisture indicator value: **2**

Reaction indicator value: **8**

Nutrient indicator value: **4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.66**

Disturbance frequency (herb layer): **1.49**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.13**

Grazing pressure: **0.44**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) Ononido-Rosmarinetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S65 Mediterranean gypsum scrub](#)

Broad habitat

Occurrence in broad habitats: **Scrub**