

# *Anthyllis hermanniae*

## Habitus and growth type

Plant height [m]: **0.4**

Life span: **Perennial**

Life form: **Chamaephyte**

Spinescence: **not spinescent**

## Flower

Flowering period: **April-July**

## Fruit, seed and dispersal

Seed mass [mg]: **1.09**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Moisture indicator value: **2.3**

Nutrient indicator value: **3**

Salinity indicator value: **0.3**

### Disturbance indicator values

Disturbance frequency: **0.35**

Disturbance frequency (herb layer): **1.32**

Disturbance severity: **0.57**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0.03**

Grazing pressure: **0.26**

Soil disturbance: **0.12**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **[FB \(ROS\) \*Ononido-Rosmarinetea\*](#), **[FQ \(GEN\) \*Carici-Genistetea lobelii\*](#)****

### EUNIS habitat

Diagnostic species of EUNIS habitats: **[R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#), [S63 Eastern garrigue](#), [S74 Central Mediterranean mountain hedgehog-heath](#)**

Constant species of EUNIS habitats: **[R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#), [S74 Central Mediterranean mountain hedgehog-heath](#)**

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree)**