Antirrhinum majus

Habitus and growth type

Plant height [m]: **0.58**
Life span: **Annual, Perennial**
Life form: **Chamaephyte, Hemicryptophyte, Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **17.2**

Flower

Flowering period: **January-November**

Fruit, seed and dispersal

Seed mass [mg]: **0.11**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**
Moisture indicator value: **3.8**
Reaction indicator value: **7.1**
Nutrient indicator value: **5.8**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.61**
Disturbance frequency (herb layer): **1.05**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.6**
Disturbance severity (herb layer): **0.45**
Mowing frequency: **0.25**
Grazing pressure: **0.21**
Soil disturbance: **0.23**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **2**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz