

# *Arabis planisiliqua*

## Habitus and growth type

Plant height [m]: **0.46**

Life span: **Biennial or short-lived**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6.9**

Moisture indicator value: **6.4**

Reaction indicator value: **7.3**

Nutrient indicator value: **3.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.18**

Disturbance frequency (herb layer): **1.42**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.14**

Grazing pressure: **0.22**

Soil disturbance: **0.16**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## **Distribution**

Continentality: **5**

Continentality amplitude: **4**