

Arabis scopoliana

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Temperature indicator value: **2**

Disturbance indicator values

Disturbance frequency: **1.12**

Disturbance frequency (herb layer): **1.12**

Disturbance severity: **0.29**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.01**

Grazing pressure: **0.25**

Soil disturbance: **0.09**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CU \(SES\) *Elyno-Seslerietea*](#), [KC \(ASP\) *Asplenietea trichomanis*](#), [KE \(THL\) *Thlaspietea rotundifolii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R45 Alpine and subalpine calcareous grassland of the Balkans and Apennines](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**