

Arenaria armerina

Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.7**

Temperature indicator value: **4**

Moisture indicator value: **2.1**

Reaction indicator value: **7.2**

Nutrient indicator value: **1.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.32**

Disturbance frequency (herb layer): **1.34**

Disturbance severity: **0.62**

Disturbance severity (herb layer): **0.2**

Mowing frequency: **0.04**

Grazing pressure: **0.32**

Soil disturbance: **0.12**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1H Iberian oromediterranean basiphilous dry grassland](#), [S73 Western Mediterranean mountain hedgehog-heath](#)

Broad habitat

Occurrence in broad habitats: **Alpine-subalpine and arctic grassland, Scrub**