Arenaria armerina

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.7**
Temperature indicator value: **4**
Moisture indicator value: **2.1**
Reaction indicator value: **7.2**
Nutrient indicator value: **1.6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.32**
Disturbance frequency (herb layer): **1.34**
Disturbance severity: **0.62**
Disturbance severity (herb layer): **0.2**
Mowing frequency: **0.04**
Grazing pressure: **0.32**
Soil disturbance: **0.12**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1H Iberian oromediterranean basiphilous dry grassland**, **S73 Western Mediterranean mountain hedgehog-heath**

Broad habitat

Occurrence in broad habitats: **Alpine-subalpine and arctic grassland, Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz