

# *Arenaria balearica*

## Habitus and growth type

Plant height [m]: **0.03**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **0.08**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **3.5**

Temperature indicator value: **6.5**

Moisture indicator value: **6.5**

Nutrient indicator value: **3.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.1**

Disturbance frequency (herb layer): **1.17**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.15**

Mowing frequency: **0.04**

Grazing pressure: **0.19**

Soil disturbance: **0.1**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [KB \(POD\) Polypodietea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T3C Taxus baccata forest](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Forest, Sparsely vegetated (incl. rock and scree)**