Arenaria controversa

Habitus and growth type

Life span: **Annual, Biennial or short-lived**
Life form: **Hemicryptophyte, Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.5**
Moisture indicator value: **3**
Reaction indicator value: **7.4**
Nutrient indicator value: **1.7**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **2.31**
Disturbance frequency (herb layer): **2.31**
Disturbance severity: **0.52**
Disturbance severity (herb layer): **0.52**
Mowing frequency: **0.39**
Grazing pressure: **0.45**
Soil disturbance: **0.36**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CJ (COR) *Koelerio-Corynephoretea canescentis***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz