

Arenaria montana

Habitus and growth type

Plant height [m]: **0.09**

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **9.19**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**

Temperature indicator value: **6.7**

Moisture indicator value: **4**

Reaction indicator value: **3.7**

Nutrient indicator value: **2.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.2**

Disturbance frequency (herb layer): **1.32**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.18**

Mowing frequency: **0.03**

Grazing pressure: **0.24**

Soil disturbance: **0.1**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [T37 Mediterranean montane Pinus sylvestris-Pinus nigra forest](#)

Broad habitat

Occurrence in broad habitats: **Forest, Sparsely vegetated (incl. rock and scree)**