

Arenaria querioides

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.28**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.7**

Temperature indicator value: **5**

Moisture indicator value: **3.1**

Reaction indicator value: **3.1**

Nutrient indicator value: **1.4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.44**

Disturbance frequency (herb layer): **1.56**

Disturbance severity: **0.3**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.11**

Grazing pressure: **0.28**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FL \(IND\) Festucetea indigestae](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1N Open Iberian supramediterranean dry](#)

acid and neutral grassland

Constant species of EUNIS habitats: **R1N Open Iberian supramediterranean dry acid and neutral grassland**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**