

Argyrolobium zanonii

Habitus and growth type

Plant height [m]: **0.12**

Life span: **Perennial**

Life form: **Chamaephyte**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **10.97**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **6.99**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.7**

Temperature indicator value: **7**

Moisture indicator value: **2.7**

Reaction indicator value: **7.3**

Nutrient indicator value: **2.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.49**
Disturbance frequency (herb layer): **1.57**
Disturbance severity: **0.62**
Disturbance severity (herb layer): **0.3**
Mowing frequency: **0.17**
Grazing pressure: **0.36**
Soil disturbance: **0.18**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [FB \(ROS\) Ononido-Rosmarinetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe](#), [S61 Western basiphilous garrigue](#)

Constant species of EUNIS habitats: [R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe](#), [S61 Western basiphilous garrigue](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**