

# *Aristolochia baetica*

## Habitus and growth type

Life span: **Perennial**

Life form: **Phanerophyte, Woody liana**

## Fruit, seed and dispersal

Seed mass [mg]: **28.16**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **6.8**

Temperature indicator value: **9**

Moisture indicator value: **3.8**

Reaction indicator value: **5.7**

Nutrient indicator value: **4.6**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.11**

Disturbance frequency (herb layer): **1.43**

Disturbance severity: **0.71**

Disturbance severity (herb layer): **0.22**

Mowing frequency: **0**

Grazing pressure: **0.27**

Soil disturbance: **0.12**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FA \(QUI\) Quercetea ilicis](#)

Broad habitat

Occurrence in broad habitats: **Forest**