Aristolochia baetica

Habitus and growth type

Life span: **Perennial**
Life form: **Phanerophyte, Woody liana**

Fruit, seed and dispersal

Seed mass [mg]: **28.16**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **6.8**
Temperature indicator value: **9**
Moisture indicator value: **3.8**
Reaction indicator value: **5.7**
Nutrient indicator value: **4.6**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.11**
Disturbance frequency (herb layer): **1.43**
Disturbance severity: **0.71**
Disturbance severity (herb layer): **0.22**
Mowing frequency: **0**
Grazing pressure: **0.27**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FA (QUI) *Quercetea ilicis***

Broad habitat

Occurrence in broad habitats: **Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz