

# *Armeria caespitosa*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **8.6**

Temperature indicator value: **3.2**

Moisture indicator value: **3.3**

Reaction indicator value: **3.5**

Nutrient indicator value: **1.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.31**

Disturbance frequency (herb layer): **1.7**

Disturbance severity: **0.33**

Disturbance severity (herb layer): **0.3**

Mowing frequency: **0.06**

Grazing pressure: **0.31**

Soil disturbance: **0.12**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [\*\*FL \(IND\) Festucetea indigestae\*\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*R1G Iberian oromediterranean siliceous dry grassland\*\*](#)

Constant species of EUNIS habitats: [\*\*R1G Iberian oromediterranean siliceous dry\*\*](#)

[grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**