Armeria leucocephala

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **7.3**
Temperature indicator value: **4.6**
Moisture indicator value: **3.6**
Reaction indicator value: **3**
Nutrient indicator value: **2.1**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.75**
Disturbance frequency (herb layer): **0.75**
Disturbance severity: **0.19**
Disturbance severity (herb layer): **0.19**
Mowing frequency: **0.02**
Grazing pressure: **0.09**
Soil disturbance: **0.03**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **U34 Mediterranean siliceous inland cliff**
Constant species of EUNIS habitats: **U34 Mediterranean siliceous inland cliff**

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz