

# *Asclepias syriaca*

## Habitus and growth type

Plant height [m]: **1.38**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **15.83**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Seed mass [mg]: **6.2**

Dispersal mode: **Anemochory, Anthropochory**

Dispersal distance class: **5, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Neophyte**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.6**

Temperature indicator value: **6.9**

Moisture indicator value: **5.1**

Reaction indicator value: **6.5**

Nutrient indicator value: **5**

Salinity indicator value: **0.3**

### Disturbance indicator values

Disturbance frequency: **1.45**

Disturbance frequency (herb layer): **1.46**

Disturbance severity: **0.62**

Disturbance severity (herb layer): **0.61**

Mowing frequency: **0.78**

Grazing pressure: **0.22**

Soil disturbance: **0.24**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) \*Koelerio-Coryneporetea canescentis\*](#), [QD \(DIG\) \*Digitario sanguinalis-Eragrostietea minoris\*](#), [QF \(ART\) \*Artemisietea vulgaris\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

## Distribution

Continentality: **6**

Continentality amplitude: **6**