

Asperula idaea

Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **8**

Moisture indicator value: **4**

Reaction indicator value: **7**

Nutrient indicator value: **4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.01**

Disturbance frequency (herb layer): **1.6**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.37**

Mowing frequency: **0.06**

Grazing pressure: **0.34**

Soil disturbance: **0.22**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FR \(DAP\) *Daphno-Festucetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S75 Eastern Mediterranean mountain hedgehog-heath](#), [U29 Eastern Mediterranean base-rich scree](#)

Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**