

Asperula tenella

Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **5.5**

Moisture indicator value: **4.5**

Disturbance indicator values

Disturbance frequency: **1.23**

Disturbance frequency (herb layer): **1.81**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.44**

Mowing frequency: **0.46**

Grazing pressure: **0.33**

Soil disturbance: **0.19**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) *Koelerio-Corynephoretea canescentis*](#), [DA \(FES\) *Festuco-Brometea*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**