

# *Asplenium seelosii*

## Habitus and growth type

Plant height [m]: **0.06**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **May-September**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Temperature indicator value: **3.7**

Moisture indicator value: **4**

Reaction indicator value: **8**

Nutrient indicator value: **1**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.6**

Disturbance frequency (herb layer): **0.66**

Disturbance severity: **0.17**

Disturbance severity (herb layer): **0.15**

Mowing frequency: **0**

Grazing pressure: **0.08**

Soil disturbance: **0.03**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [KC \(ASP\) \*Asplenietea trichomanis\*](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Spring, Sparsely vegetated (incl. rock and scree)**

## **Distribution**

Continentality: **4**

Continentality amplitude: **2**