

Astragalus exscapus

Habitus and growth type

Plant height [m]: **0.14**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **5.91**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.4**

Temperature indicator value: **5.9**

Moisture indicator value: **2.9**

Reaction indicator value: **8.8**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.1**

Disturbance frequency (herb layer): **1.58**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.53**

Grazing pressure: **0.33**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [BD \(PYR\) *Pyrolo-Pinetetea sylvestris*](#),
[DA \(FES\) *Festuco-Brometea*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **8**

Continentality amplitude: **4**