

# *Astragalus monspessulanus*

## Habitus and growth type

Plant height [m]: **0.18**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **13.97**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **2.05**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.6**

Temperature indicator value: **6.9**

Moisture indicator value: **3.1**

Reaction indicator value: **8**

Nutrient indicator value: **1.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.48**

Disturbance frequency (herb layer): **1.57**

Disturbance severity: **0.57**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.24**

Grazing pressure: **0.32**

Soil disturbance: **0.16**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*FB \(ROS\) Ononido-Rosmarinetea\*\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe\*\*](#)

Constant species of EUNIS habitats: [\*\*R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe\*\*](#), [\*\*S61 Western basiphilous garrigue\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**