

Astragalus tragacantha

Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte**

Fruit, seed and dispersal

Seed mass [mg]: **3.3**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **9**

Moisture indicator value: **5**

Reaction indicator value: **6**

Nutrient indicator value: **5**

Salinity indicator value: **6**

Disturbance indicator values

Disturbance frequency: **0.76**

Disturbance frequency (herb layer): **0.87**

Disturbance severity: **0.38**

Disturbance severity (herb layer): **0.3**

Mowing frequency: **0.11**

Grazing pressure: **0.21**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [JB \(CRI\) Crithmo-Staticetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S71 Western Mediterranean spiny heath](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff**