

Avenula pubescens

Habitus and growth type

Plant height [m]: **0.63**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **18.42**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **1.85**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.5**

Temperature indicator value: **4.9**

Moisture indicator value: **3.5**

Reaction indicator value: **6.1**

Nutrient indicator value: **3.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.66**

Disturbance frequency (herb layer): **1.98**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.48**

Mowing frequency: **1.06**

Grazing pressure: **0.19**

Soil disturbance: **0.29**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R22 Low and medium altitude hay meadow](#),
[R23 Mountain hay meadow](#)

Constant species of EUNIS habitats: [R1A Semi-dry perennial calcareous grassland \(meadow steppe\)](#), [R22 Low and medium altitude hay meadow](#), [R23 Mountain hay meadow](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: **6**