

# ***Barbarea stricta***

## **Habitus and growth type**

Plant height [m]: **0.6**

Life span: **Biennial or short-lived, Perennial**

Life form: **Hemicryptophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **43.36**

## **Flower**

Flowering period: **March-June**

## **Fruit, seed and dispersal**

Seed mass [mg]: **0.32**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **7.9**

Moisture indicator value: **6.8**

Reaction indicator value: **7**

Nutrient indicator value: **7**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **0.78**

Disturbance frequency (herb layer): **1.42**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **0.44**

Grazing pressure: **0.18**

Soil disturbance: **0.21**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [QG \(EPI\) Epilobietea angustifolii](#)

### Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Synanthropic**

## Distribution

Continentality: **6**

Continentality amplitude: **6**