

Bellardia trixago

Habitus and growth type

Plant height [m]: **0.26**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **12.22**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **8.5**

Moisture indicator value: **3**

Reaction indicator value: **5.7**

Nutrient indicator value: **3.7**

Salinity indicator value: **0.5**

Disturbance indicator values

Disturbance frequency: **1.31**

Disturbance frequency (herb layer): **1.86**

Disturbance severity: **0.72**

Disturbance severity (herb layer): **0.65**

Mowing frequency: **0.51**

Grazing pressure: **0.3**

Soil disturbance: **0.35**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) *Chenopodietea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S72 Eastern Mediterranean spiny heath \(phrygana\)](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Synanthropic**