

Berula erecta

Habitus and growth type

Plant height [m]: **0.58**

Life span: **Perennial**

Life form: **Hemicryptophyte, Hydrophyte**

Leaf

Specific leaf area [mm²/mg]: **33.66**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.63**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **5.5**

Moisture indicator value: **9.9**

Reaction indicator value: **7.5**

Nutrient indicator value: **6.1**

Salinity indicator value: **0.6**

Disturbance indicator values

Disturbance frequency: **1.33**

Disturbance frequency (herb layer): **1.72**
Disturbance severity: **0.31**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.2**
Grazing pressure: **0.14**
Soil disturbance: **0.08**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OD \(PHR\) *Phragmito-Magnocaricetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q52 *Small-helophyte bed*](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland**

Distribution

Continentality amplitude: 7