

# ***Betula pubescens***

## **Habitus and growth type**

Plant height [m]: **19.19**

Life span: **Perennial**

Life form: **Phanerophyte, Tree**

## **Leaf**

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

Specific leaf area [mm<sup>2</sup>/mg]: **14**

## **Flower**

Flowering period: **March-May**

## **Fruit, seed and dispersal**

Seed mass [mg]: **0.33**

Dispersal mode: **Dyszoochory**

Dispersal distance class: **6**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **4.2**

Reaction indicator value: **3**

Nutrient indicator value: **2.9**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.22**  
Disturbance frequency (herb layer): **1.22**  
Disturbance severity: **0.62**  
Disturbance severity (herb layer): **0.16**  
Mowing frequency: **0.05**  
Grazing pressure: **0.18**  
Soil disturbance: **0.16**

## Habitat and sociology

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T16 Broadleaved mire forest on acid peat](#), [T1C Temperate and boreal mountain Betula and Populus tremula forest on mineral soils](#), [T3F Dark taiga](#), [T3G Pinus sylvestris light taiga](#), [T3H Larix light taiga](#), [T3J Pinus and Larix mire forest](#), [T3K Picea mire forest](#)

Constant species of EUNIS habitats: [N1D Atlantic and Baltic broad-leaved coastal dune forest](#), [N1F Baltic coniferous coastal dune forest](#), [Q11 Raised bog](#), [Q22 Poor fen](#), [Q42 Extremely rich moss-sedge fen](#), [S41 Wet heath](#), [S92 Salix fen scrub](#), [T15 Broadleaved swamp forest on non-acid peat](#), [T16 Broadleaved mire forest on acid peat](#), [T1B Acidophilous Quercus forest](#), [T1C Temperate and boreal mountain Betula and Populus tremula forest on mineral soils](#), [T27 Ilex aquifolium forest](#), [T3F Dark taiga](#), [T3G Pinus sylvestris light taiga](#), [T3H Larix light taiga](#), [T3J Pinus and Larix mire forest](#), [T3K Picea mire forest](#), [U61 Subarctic volcanic field](#)

Dominant species of EUNIS habitats: [T16 Broadleaved mire forest on acid peat](#), [T1C Temperate and boreal mountain Betula and Populus tremula forest on mineral soils](#)

### Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Mire, Grassland (non-alpine, non-saline), Heathland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**