Biarum tenuifolium

Habitus and growth type

Plant height [m]: **0.3**
Life span: **Perennial**
Life form: **Geophyte**

Flower

Flowering period: **January-June, September-December**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.8**
Temperature indicator value: **9**
Moisture indicator value: **4**
Nutrient indicator value: **5.8**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.25**
Disturbance frequency (herb layer): **1.46**
Disturbance severity: **0.72**
Disturbance severity (herb layer): **0.38**
Mowing frequency: **0.12**
Grazing pressure: **0.31**
Soil disturbance: **0.15**

Habitat and sociology

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz