

Blitum virgatum

Habitus and growth type

Plant height [m]: **0.46**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.56**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.1**

Temperature indicator value: **5.8**

Moisture indicator value: **4.2**

Reaction indicator value: **6.3**

Nutrient indicator value: **7.4**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.43**

Disturbance frequency (herb layer): **1.61**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.69**

Mowing frequency: **0.61**

Grazing pressure: **0.16**

Soil disturbance: **0.43**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QF \(ART\) Artemisietea vulgaris](#)

Broad habitat

Occurrence in broad habitats: **Synanthropic**

Distribution

Continentality: **7**

Continentality amplitude: **6**