

Bupleurum semicompositum

Habitus and growth type

Plant height [m]: **0.18**

Life span: **Annual**

Life form: **Therophyte**

Spinescence: **not spinescent**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.2**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **11**

Moisture indicator value: **4.5**

Reaction indicator value: **9**

Nutrient indicator value: **3**

Salinity indicator value: **6**

Disturbance indicator values

Disturbance frequency: **1.4**

Disturbance frequency (herb layer): **1.51**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.63**

Mowing frequency: **0.13**

Grazing pressure: **0.32**

Soil disturbance: **0.31**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FJ \(TRA\) Stipo-Trachynietea distachyae](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R61 Mediterranean inland salt steppe](#), [S66 Mediterranean halo-nitrophilous scrub](#)

Constant species of EUNIS habitats: [S66 Mediterranean halo-nitrophilous scrub](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**