Calicotome villosa

Habitus and growth type

Plant height [m]: **1.5**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**
Spinescence: **spinescent**

Leaf

Specific leaf area [mm^2/mg]: **13.9**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **7.16**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**
Temperature indicator value: **9**
Moisture indicator value: **3**
Reaction indicator value: **6**
Nutrient indicator value: **3**
Salinity indicator value: **0**

Disturbance indicator values

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency: **0.36**
Disturbance frequency (herb layer): **1.41**
Disturbance severity: **0.71**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.02**
Grazing pressure: **0.31**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FA (QUI) *Quercetea ilicis***, **FB (ROS) *Ononido-Rosmarinetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S51 Mediterranean maquis and arborescent matorral**, **S54 Thermomediterranean arid scrub**, **S63 Eastern garrigue**, **S71 Western Mediterranean spiny heath**, **S72 Eastern Mediterranean spiny heath (phrygana)**, **T24 Olea europaea-Ceratonia siliqua forest**
Constant species of EUNIS habitats: **S51 Mediterranean maquis and arborescent matorral**, **S54 Thermomediterranean arid scrub**, **S63 Eastern garrigue**, **S71 Western Mediterranean spiny heath**, **S72 Eastern Mediterranean spiny heath (phrygana)**, **T24 Olea europaea-Ceratonia siliqua forest**
Dominant species of EUNIS habitats: **S51 Mediterranean maquis and arborescent matorral**

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz