

# *Callitriche brutia*

## Habitus and growth type

Plant height [m]: **0.17**

Life span: **Annual, Perennial**

Life form: **Hydrophyte, Therophyte**

## Flower

Flowering period: **April-October**

## Fruit, seed and dispersal

Seed mass [mg]: **0.66**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **6.8**

Moisture indicator value: **10.3**

Reaction indicator value: **5**

Nutrient indicator value: **4.5**

Salinity indicator value: **0.5**

### Disturbance indicator values

Disturbance frequency: **1.84**

Disturbance frequency (herb layer): **1.84**

Disturbance severity: **0.28**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.01**

Grazing pressure: **0.03**

Soil disturbance: **0.08**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [NB \(POT\) Potamogetonetea](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland**

## **Distribution**

Continentality: **2**

Continentality amplitude: **2**