

Caltha palustris

Habitus and growth type

Plant height [m]: **0.33**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **27.39**

Flower

Flowering period: **February-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.98**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Temperature indicator value: **4.6**

Moisture indicator value: **9**

Reaction indicator value: **4.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.77**

Disturbance frequency (herb layer): **1.58**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.41**

Mowing frequency: **0.41**

Grazing pressure: **0.18**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**CM \(MOL\) Molinio-Arrhenatheretea**](#),
[**OA \(MON\) Montio-Cardaminetea**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**R35 Moist or wet mesotrophic to eutrophic hay meadow**](#), [**T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils**](#)

Constant species of EUNIS habitats: [**P2N Spring**](#), [**Q23 Relict mire of Mediterranean mountains**](#), [**Q24 Intermediate fen and soft-water spring mire**](#), [**Q41 Alkaline, calcareous, carbonate-rich small-sedge spring fen**](#), [**Q42 Extremely rich moss-sedge fen**](#), [**Q53 Tall-sedge bed**](#), [**R35 Moist or wet mesotrophic to eutrophic hay meadow**](#), [**R55 Lowland moist or wet tall-herb and fern fringe**](#), [**T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils**](#), [**T13 Temperate hardwood riparian forest**](#), [**T15 Broadleaved swamp forest on non-acid peat**](#), [**U3D Wet inland cliff**](#)

Dominant species of EUNIS habitats: [**P2N Spring**](#)

Broad habitat

Occurrence in broad habitats: **Spring, Wetland, Mire, Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: **8**