

# *Camelina microcarpa*

## Habitus and growth type

Plant height [m]: **0.35**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **19.99**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **0.31**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.4**

Moisture indicator value: **3.9**

Reaction indicator value: **7.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.64**

Disturbance frequency (herb layer): **1.94**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.68**

Mowing frequency: **0.59**

Grazing pressure: **0.24**

Soil disturbance: **0.5**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [QA \(PAR\) Papaveretea rhoeadis](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [S67 Aralo-Caspian semi-desert](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**

## Distribution

Continentality: **7**

Continentality amplitude: **5**