

Campanula glomerata

Habitus and growth type

Plant height [m]: **0.36**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **30.72**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.13**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **5.6**

Moisture indicator value: **4.1**

Reaction indicator value: **7**

Nutrient indicator value: **3.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.83**

Disturbance frequency (herb layer): **1.75**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.42**
Mowing frequency: **0.61**
Grazing pressure: **0.24**
Soil disturbance: **0.17**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [BC \(BRA\) *Brachypodio pinnati-Betuletea pendulae*](#), [CB \(PUB\) *Quercetea pubescentis*](#), [CM \(MOL\) *Molinio-Arrhenatheretea*](#), [CU \(SES\) *Elyno-Seslerietea*](#), [DA \(FES\) *Festuco-Brometea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T3H *Larix light taiga*](#)

Constant species of EUNIS habitats: [T3H *Larix light taiga*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: 7

Continentality amplitude: 7