

Campanula radicata

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.4**

Temperature indicator value: **5.5**

Moisture indicator value: **5.8**

Reaction indicator value: **6.3**

Nutrient indicator value: **3.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.31**

Disturbance frequency (herb layer): **2.31**

Disturbance severity: **0.33**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.02**

Grazing pressure: **0.34**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **[FO \(ANA\) *Trifolium anatolici-*](#)**

[Polygonetia arenastri](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: **[R1K Balkan and Anatolian oromediterranean dry grassland](#)**

Constant species of EUNIS habitats: **[R1K Balkan and Anatolian oromediterranean dry grassland](#)**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**