

Campanula trachelium

Habitus and growth type

Plant height [m]: **0.61**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]:

Flower

Flowering period: **May-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.18**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4.4**

Temperature indicator value: **5.4**

Moisture indicator value: **5.5**

Reaction indicator value: **7.8**

Nutrient indicator value: **7.1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.07**

Disturbance frequency (herb layer): **1.32**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.16**
Mowing frequency: **0.03**
Grazing pressure: **0.19**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) *Carpino-Fagetea sylvaticae*](#)

EUNIS habitat

Constant species of EUNIS habitats: [S37 *Corylus avellana* scrub](#), [T17 *Fagus* forest on non-acid soils](#), [T19 Temperate and submediterranean thermophilous deciduous forest](#), [T1E *Carpinus* and *Quercus* mesic deciduous forest](#), [T1F Ravine forest](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Synanthropic**

Distribution

Continentality amplitude: **5**