

Cardamine heptaphylla

Habitus and growth type

Plant height [m]: **0.42**

Life span: **Perennial**

Life form: **Geophyte**

Leaf

Specific leaf area [mm²/mg]: **38.7**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **6.39**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **3**

Temperature indicator value: **4.5**

Moisture indicator value: **5**

Reaction indicator value: **7.7**

Nutrient indicator value: **6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.03**

Disturbance frequency (herb layer): **1.3**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.12**
Mowing frequency: **0.01**
Grazing pressure: **0.18**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) *Carpino-Fagetea sylvaticae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T32 Temperate mountain *Abies* forest](#)

Constant species of EUNIS habitats: [T32 Temperate mountain *Abies* forest](#)

Broad habitat

Occurrence in broad habitats: **Forest**

Distribution

Continentality: **3**

Continentality amplitude: **1**