Cardamine raphanifolia

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **4**
Moisture indicator value: **9**
Reaction indicator value: **4.5**
Nutrient indicator value: **4.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.36**
Disturbance frequency (herb layer): **1.47**
Disturbance severity: **0.56**
Disturbance severity (herb layer): **0.3**
Mowing frequency: **0.06**
Grazing pressure: **0.17**
Soil disturbance: **0.12**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OA (MON) *Montio-Cardaminetea***

Broad habitat

Occurrence in broad habitats: **Spring**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz