

# *Carex brizoides*

## Habitus and growth type

Plant height [m]: **0.47**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **26.35**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **0.39**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **5.8**

Temperature indicator value: **5.5**

Reaction indicator value: **3.6**

Nutrient indicator value: **3.1**

Salinity indicator value: **0.1**

### Disturbance indicator values

Disturbance frequency: **0.24**

Disturbance frequency (herb layer): **1.46**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.14**

Grazing pressure: **0.19**

Soil disturbance: **0.13**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#),  
[HA \(POP\) Alno glutinosae-Populetea albae](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils](#), [T13 Temperate hardwood riparian forest](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Synanthropic**

## Distribution

Continentality: **5**

Continentality amplitude: **3**