Carex camposii

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Disturbance indicator values

Disturbance frequency: **0.96**
Disturbance frequency (herb layer): **2.04**
Disturbance severity: **0.49**
Disturbance severity (herb layer): **0.4**
Mowing frequency: **0.54**
Grazing pressure: **0.46**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OD (PHR) *Phragmito-Magnocaricetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R24 Iberian summer pasture (vallicar)**

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz