

Carex capillaris

Habitus and growth type

Plant height [m]: **0.19**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **13.23**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.55**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **1.7**

Reaction indicator value: **7.6**

Nutrient indicator value: **1.9**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.35**

Disturbance frequency (herb layer): **1.54**

Disturbance severity: **0.27**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.12**

Grazing pressure: **0.19**

Soil disturbance: **0.07**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [AB \(KOB\) *Carici rupestris-Kobresietea bellardii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q45 Arctic-alpine rich fen](#), [U35 Boreal and Arctic base-rich inland cliff](#)

Constant species of EUNIS habitats: [Q45 Arctic-alpine rich fen](#), [U25 Boreal and Arctic base-rich scree and block field](#), [U35 Boreal and Arctic base-rich inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Spring, Wetland, Mire, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**

Continentality amplitude: **6**