

Carex filiformis

Habitus and growth type

Plant height [m]: **0.36**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

Leaf

Specific leaf area [mm²/mg]: **18.82**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.1**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.9**

Temperature indicator value: **5.5**

Moisture indicator value: **6.6**

Reaction indicator value: **8.3**

Nutrient indicator value: **3.8**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.09**

Disturbance frequency (herb layer): **1.87**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.94**

Grazing pressure: **0.24**

Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) *Molinio-Arrhenatheretea*](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Mire, Grassland (non-alpine, non-saline), Saline vegetation**

Distribution

Continentality: **6**

Continentality amplitude: **5**